

Generalized functions valued in a smooth manifold

Michael Kunzinger *

Department of Mathematics, University of Vienna
Strudlhofg. 4, A-1090 Wien, Austria

Abstract

Based on Colombeau's theory of algebras of generalized functions we introduce the concepts of generalized functions taking values in differentiable manifolds as well as of generalized vector bundle homomorphisms. We study their basic properties, in particular with respect to some new point value concepts for generalized functions and indicate applications of the resulting theory in general relativity.

Mathematics Subject Classification (2000): Primary: 46T30; secondary: 46F30, 53B20.

Keywords Algebras of generalized functions, Colombeau algebras, generalized functions on manifolds.

1 Introduction

An increasing number of applications of Colombeau's theory of algebras of generalized functions ([2, 3, 16]) to questions of a primarily geometric nature, in particular in general relativity (cf. [19] and the literature cited therein for a survey) have resulted in a certain restructuring of the theory itself. For its full version, this task has been pursued in [6, 7]. For the special version of the theory, which will also provide the framework of the present article, the program of "geometrizing" the construction has been initiated in [5, 12]. In particular, in [12] generalized sections of vector bundles were introduced, a point value description of Colombeau functions on manifolds was given and a number of consistency results with linear distributional geometry in the sense of [15] were derived.

When analyzing singularities in a global context by means of algebras of generalized functions one inevitably encounters situations where a concept of generalized mappings *valued* in differentiable manifolds is called for. As examples we mention the flow of a generalized vector field on a manifold or the

*Electronic mail: Michael.Kunzinger@univie.ac.at

notion of geodesic of a distributional spacetime metric. Both notions have been treated in the Colombeau framework, however, due to the lack of a global notion of generalized mappings between manifolds this treatment was confined strictly to the local context (cf. e.g., [4, 9, 10, 11, 19, 20]).

In the present paper we introduce the space $\mathcal{G}[X, Y]$ of generalized mappings defined on the manifold X and taking values in the manifold Y , adapted to the requirements of applying algebras of generalized functions to global analysis in the presence of singularities (Section 2). The basic idea underlying the construction is to consider nets of smooth maps $u_\varepsilon : X \rightarrow Y$ parametrized by a regularization parameter $\varepsilon > 0$ such that for any compact set K in X and small ε the values of all u_ε on K remain in a compact subset of Y . In addition we will require a moderateness condition on the derivatives of the u_ε (growth on compact sets bounded by certain inverse powers of ε). This growth restriction has to be formulated in local charts. An equivalence relation on such nets will be introduced by employing Riemannian metrics on the manifolds. However, the definition will turn out to be independent of the chosen metric. Switching between coordinate descriptions and global formulations in terms of a (but independent of the particular) Riemannian metric will be a reoccurring pattern throughout this paper.

Typical singularities that can be modelled by elements of $\mathcal{G}[X, Y]$ are jump discontinuities. Singularities of a less tame nature (e.g., delta-type singularities) of course have to be handled when differentiating elements of $\mathcal{G}[X, Y]$. In order to accommodate such less pleasant creatures, i.e., to develop a suitable notion of tangent map for elements of $\mathcal{G}[X, Y]$ we therefore have to consider generalized vector bundle homomorphisms (Section 3) which will be allowed to behave more singularly in their vector components. In this context a new notion of generalized points in vector bundles is introduced that allows to characterize generalized sections of vector bundles. Finally, we show that the above “generalized vector bundle points” are of independent value in that they allow a direct transfer of the “pointwise” nature of smooth tensor fields to the generalized context.

In the remainder of the introduction we fix some notations from global analysis and recall those notions and results from [12] which will be needed in our presentation.

X, Y will denote smooth paracompact Hausdorff manifolds of dimensions n and m , respectively. Vector bundles with base space X will be denoted by (E, X, π) (or (E, X, π_X)) and for a chart (V, φ) in X , a vector bundle chart (V, Φ) over φ will be written in the form ($\mathbb{K} = \mathbb{R}$ for real resp. $\mathbb{K} = \mathbb{C}$ for complex vector bundles)

$$\begin{aligned} \Phi : \pi^{-1}(V) &\rightarrow \varphi(V) \times \mathbb{K}^{n'} \\ z &\rightarrow (\varphi(p), \varphi^1(z), \dots, \varphi^{n'}(z)) \equiv (\varphi(p), \boldsymbol{\varphi}(z)), \end{aligned} \quad (1)$$

where $p = \pi(z)$ and the typical fiber is $\mathbb{K}^{n'}$. Let $(V_\alpha, \Phi_\alpha)_\alpha$ denote a vector bundle atlas; then we write $\Phi_\alpha \circ \Phi_\beta^{-1}(y, w) = (\varphi_{\alpha\beta}(y), \boldsymbol{\varphi}_{\alpha\beta}(y)w)$, where $\varphi_{\alpha\beta} := \varphi_\alpha \circ \varphi_\beta^{-1}$ is the change of chart on the base and $\boldsymbol{\varphi}_{\alpha\beta} : \varphi_\beta(V_\alpha \cap V_\beta) \rightarrow \text{GL}(n', \mathbb{K})$

denotes the transition functions.

For vector bundles (E, X, π_X) , (F, Y, π_Y) we denote by $\text{Hom}(E, F)$ the space of smooth vector bundle homomorphisms from E to F . If $f \in \text{Hom}(E, F)$ we denote by $\underline{f} : X \rightarrow Y$ the smooth map induced by f on the base manifolds. Thus $\pi_Y \circ f = \underline{f} \circ \pi_X$. For vector bundle charts (V, Φ) of E and (W, Ψ) of F we write the local vector bundle homomorphism $f_{\Psi\Phi} := \Psi \circ f \circ \Phi^{-1} : \varphi(V \cap \underline{f}^{-1}(W)) \times \mathbb{K}^{n'} \rightarrow \psi(W) \times \mathbb{K}^{m'}$ in the form

$$f_{\Psi\Phi}(x, \xi) = (f_{\Psi\Phi}^{(1)}(x), f_{\Psi\Phi}^{(2)}(x) \cdot \xi). \quad (2)$$

The space of $\mathcal{C}^\infty$ -sections of (E, X, π) is denoted by $\Gamma(X, E)$. As a special case we mention $\Gamma(X, T_s^r(X)) = \mathcal{T}_s^r(X)$, the space of (r, s) -tensor fields over X .

Turning now to notations from Colombeau theory we set $I = (0, 1]$, let $\mathcal{P}(X)$ be the space of linear differential operators on X and define

$$\begin{aligned} \mathcal{E}(X) &:= \mathcal{C}^\infty(X)^I \\ \mathcal{E}_M(X) &:= \{(u_\varepsilon)_\varepsilon \in \mathcal{E}(X) : \forall K \subset\subset X, \forall P \in \mathcal{P}(X) \exists N \in \mathbb{N} : \\ &\quad \sup_{p \in K} |Pu_\varepsilon(p)| = O(\varepsilon^{-N})\} \\ \mathcal{N}(X) &:= \{(u_\varepsilon)_\varepsilon \in \mathcal{E}_M(X) : \forall K \subset\subset X, \forall q \in \mathbb{N}_0 : \sup_{p \in K} |u_\varepsilon(p)| = O(\varepsilon^q)\}. \end{aligned}$$

$\mathcal{G}(X) := \mathcal{E}_M(X)/\mathcal{N}(X)$ is called the (special) Colombeau algebra on X . $u \in \mathcal{G}(X)$ if and only if $u \circ \psi_\alpha \in \mathcal{G}(\psi_\alpha(V_\alpha))$ (the local Colombeau algebra on $\psi_\alpha(V_\alpha)$) for all charts (V_α, ψ_α) . Hence $\mathcal{G}(X)$ can be identified with the space of families $(u_\alpha)_\alpha$ with $u_\alpha \in \mathcal{G}(\psi_\alpha(V_\alpha))$ satisfying $u_\alpha|_{\psi_\alpha(V_\alpha \cap V_\beta)} = u_\beta|_{\psi_\beta(V_\alpha \cap V_\beta)} \circ \psi_\beta \circ \psi_\alpha^{-1}$ for all α, β with $V_\alpha \cap V_\beta \neq \emptyset$. $\mathcal{C}^\infty(X)$ is a subalgebra of the differential algebra $\mathcal{G}(X)$ and there exist injective sheaf morphisms embedding $\mathcal{D}'(X)$, the space of Schwartz distributions on X , into $\mathcal{G}(X)$.

A net $(p_\varepsilon)_\varepsilon \in X^I$ is called compactly supported if there exists some $K \subset\subset X$ and some $\varepsilon_0 > 0$ such that $p_\varepsilon \in K$ for $\varepsilon < \varepsilon_0$. Two compactly supported points $(p_\varepsilon)_\varepsilon, (q_\varepsilon)_\varepsilon$ are called equivalent, $(p_\varepsilon)_\varepsilon \sim (q_\varepsilon)_\varepsilon$, if $d_h(p_\varepsilon, q_\varepsilon) = O(\varepsilon^m)$ for each $m > 0$ for the distance function d_h induced on X by a Riemannian metric h on X . This notion is independent of the particular h . The quotient space $\tilde{X}_c$ of the set of compactly supported points modulo this equivalence relation is called the space of compactly supported generalized points on X . Its elements are written as $\tilde{p} = \text{cl}[(p_\varepsilon)_\varepsilon]$. For $u \in \mathcal{G}(X)$, $u(\tilde{p}) := \text{cl}[(u_\varepsilon(p_\varepsilon))_\varepsilon]$ yields a well-defined element of $\mathcal{K}$, the space of generalized numbers (see [17]). Any $u \in \mathcal{G}(X)$ is uniquely determined by its point values on $\tilde{X}_c$ ([12], Th. 1).

Generalized sections of E are defined as elements of $\Gamma_{\mathcal{G}}(X, E)$, where (with $\mathcal{P}(X, E)$ the space of linear differential operators on $\Gamma(X, E)$)

$$\begin{aligned} \Gamma_{\mathcal{E}_M}(X, E) &:= \{(s_\varepsilon)_{\varepsilon \in I} \in \Gamma(X, E)^I : \forall P \in \mathcal{P}(X, E) \forall K \subset\subset X \exists N \in \mathbb{N} : \\ &\quad \sup_{p \in K} \|Pu_\varepsilon(p)\|_h = O(\varepsilon^{-N})\} \\ \Gamma_{\mathcal{N}}(X, E) &:= \{(s_\varepsilon)_{\varepsilon \in I} \in \Gamma_{\mathcal{E}_M}(X, E) : \forall K \subset\subset X \forall m \in \mathbb{N} : \end{aligned}$$

$$\sup_{p \in K} \|u_\varepsilon(p)\|_h = O(\varepsilon^m)\}, \text{ and, finally}$$

$$\Gamma_{\mathcal{G}}(X, E) := \Gamma_{\mathcal{E}_M}(X, E) / \Gamma_{\mathcal{N}}(X, E).$$

Here $\|\cdot\|_h$ denotes the norm induced on the fibers of E by any Riemannian metric h on E .

The $\mathcal{C}^\infty(X)$ -module $\Gamma_{\mathcal{G}}(X, E)$ is algebraically characterized by (see [12], Th. 4)

$$\Gamma_{\mathcal{G}}(X, E) = \mathcal{G}(X) \otimes_{\mathcal{C}^\infty(X)} \Gamma(X, E), \quad (3)$$

Finally, generalized tensor fields are introduced as generalized sections of the bundle $T_s^r(X)$. The space of generalized (r, s) -tensor fields is denoted by $\mathcal{G}_s^r(X)$. Algebraically, the following characterizations have been established in [12], Th. 7.

$$\begin{aligned} \mathcal{G}_s^r(X) &\cong L_{\mathcal{C}^\infty(X)}(\mathfrak{X}^*(X)^r, \mathfrak{X}(X)^s; \mathcal{G}(X)) \\ \mathcal{G}_s^r(X) &\cong L_{\mathcal{G}(X)}(\mathcal{G}_1^0(X)^r, \mathcal{G}_0^1(X)^s; \mathcal{G}(X)) \\ \mathcal{G}_s^r(X) &\cong \mathcal{G}(X) \otimes_{\mathcal{C}^\infty(X)} \mathcal{T}_s^r(X). \end{aligned}$$

Here, the first line gives a $\mathcal{C}^\infty(X)$ -module isomorphism, the second a $\mathcal{G}(X)$ -isomorphism and the third line is valid in both senses.

2 Generalized mappings valued in a manifold

When generalizing the space of Colombeau generalized functions on the open set Ω taking “values” in $\mathbb{R}^m$ to the manifold setting, the first problem, namely passing from Ω to X has been dealt with successfully in [5, 12]. Any attempt to replace the range space $\mathbb{R}^m$ by some manifold Y in a straightforward manner, however, leads into serious troubles due to the linear structure of $\mathbb{R}^m$ being used in an essential way when defining moderateness and negligibility, respectively.

The first idea which comes to mind, of course, is to take a local point of view and to try to formulate moderateness and negligibility in terms of coordinate representations of representatives $(u_\varepsilon)_\varepsilon$. Yet, transforming the functions $u_\varepsilon : \mathbb{R} \rightarrow (0, 2)$, $u_\varepsilon(x) := \varepsilon$ by the diffeomorphism $\psi : (0, 2) \rightarrow (e^{1/2}, \infty)$, $\psi(y) := e^{1/y}$ on the range space shows that this naïve approach is bound to fail since it does not even lead to a well-defined notion of moderateness.

A solution to this problem can be based on generalizing the space $\mathcal{G}[\Omega, \Omega']$ introduced in [1], 7.3 (denoted there by $\mathcal{G}_*(\Omega, \Omega')$) to the manifold setting. We shortly recall its definition. Let

$$\mathcal{E}[\Omega, \Omega'] = \{(u_\varepsilon)_\varepsilon \in \mathcal{C}^\infty(\Omega, \Omega')^I \mid \forall K \subset\subset \Omega \exists \varepsilon_0 \exists K' \subset\subset \Omega' \forall \varepsilon < \varepsilon_0 u_\varepsilon(K) \subseteq K'\}$$

and denote by $\mathcal{E}_M[\Omega, \Omega']$ the subset of moderate elements of $\mathcal{E}[\Omega, \Omega']$, i.e., we set $\mathcal{E}_M[\Omega, \Omega'] = \mathcal{E}[\Omega, \Omega'] \cap \mathcal{E}_M(\Omega)^m$. Then $\mathcal{G}[\Omega, \Omega'] = \mathcal{E}_M[\Omega, \Omega'] / (\mathcal{N}(\Omega)^m)$. In [1], $\mathcal{G}[\Omega, \Omega']$ is called the space of Colombeau generalized functions *valued* in Ω' . Since in this terminology the notion “generalized function valued in $\mathbb{R}$ ” would

be ambiguous (referring to both $\mathcal{G}(\Omega)$ and $\mathcal{G}[\Omega, \mathbb{R}]$) we prefer to adopt the term “c(ompactly)-bounded” for the generalized functions under consideration.

In defining $\mathcal{E}_M[\Omega, \Omega']$ the functions themselves (i.e., derivatives of order zero) are controlled by a purely topological condition while, as it will turn out, for the equivalence relation corresponding to factoring out the null ideal, one of the conditions on the zero-th derivative can be formulated on a manifold by means of a distance function generated by some Riemannian metric. Thus in both cases, we are led to conditions of intrinsic character on the manifold. Concerning general derivatives (of order at least 1 in the case of moderateness), the usual estimates can be transferred to the manifold setting using local language, due to the chain rule.

There is one technicality to be observed in the latter process: Fixing a compact subset K on the manifold X and studying the behavior of $(u_\varepsilon)_\varepsilon$ on X by means of local charts (V, φ) and (W, ψ) in X resp. Y , it may well happen that K contains boundary points of V and/or $u_\varepsilon^{-1}(W)$. Towards the boundary, however, the functions constituting a chart may tend to infinity arbitrarily fast. Therefore, we have to restrict our attention to compact subsets L and L' of V resp. W when controlling growth of derivatives in local terms.

Summarizing, our definitions eventually will be of partly geometric and partly local character. The following proposition suggests the correct way of extracting relevant information on the moderateness of $(u_\varepsilon)_\varepsilon$ from its local expressions in charts.

2.1 Proposition. *Let $(u_\varepsilon)_\varepsilon \in \mathcal{C}^\infty(\Omega, \Omega')^I$. The following are equivalent:*

- (a) $(u_\varepsilon)_\varepsilon \in \mathcal{E}_M[\Omega, \Omega']$.
- (b) (i) $\forall K \subset\subset \Omega \exists \varepsilon_0 > 0 \exists K' \subset\subset \Omega' \forall \varepsilon < \varepsilon_0 : u_\varepsilon(K) \subseteq K'$.
(ii) $\forall k \in \mathbb{N}$, for each chart (V, φ) in Ω , each chart (W, ψ) in Ω' , each $L \subset\subset V$ and each $L' \subset\subset W$ there exists $N \in \mathbb{N}$ with

$$\sup_{x \in L \cap u_\varepsilon^{-1}(L')} \|D^{(k)}(\psi \circ u_\varepsilon \circ \varphi^{-1})(\varphi(x))\| = O(\varepsilon^{-N}).$$

Remark. (i) Here $D^{(k)}$ denotes the operation of forming the total derivative of order k and $\| \cdot \|$ is any norm on the respective space of multilinear maps (the condition is clearly independent of the choice of $\| \cdot \|$).

(ii) Explicitly, the estimate in (b)(ii) means:

$$\begin{aligned} \exists \varepsilon_1 > 0 \exists C > 0 \forall \varepsilon < \varepsilon_1 \forall x \in L \cap u_\varepsilon^{-1}(L') : \\ \|D^{(k)}(\psi \circ u_\varepsilon \circ \varphi^{-1})(\varphi(x))\| \leq C\varepsilon^{-N}. \end{aligned}$$

Therefore, let us agree to treat a supremum over the empty set as having the value zero in the present context.

Proof. (a) $\Rightarrow$ (b): (i) holds by definition. Concerning (ii), let $k \in \mathbb{N}$ and choose $N \in \mathbb{N}$, $\varepsilon_1 > 0$ such that $\|D^{(j)}u_\varepsilon(x)\| \leq \varepsilon^{-N}$ for $1 \leq j \leq k$, $\varepsilon < \varepsilon_1$ and all $x \in L$. Then by the chain rule, for $x \in L$ with $u_\varepsilon(x) \in L'$, $\|D^{(k)}(\psi \circ u_\varepsilon \circ \varphi^{-1})(\varphi(x))\|$

can be estimated by $C\varepsilon^{-N}$, where C depends only on $\sup_{y \in L'} \|D^{(j_1)}\psi(y)\|$ and $\sup_{z \in \varphi(L)} \|D^{(j_2)}(\varphi^{-1})(z)\|$ ($1 \leq j_1, j_2 \leq k$), so the claim follows.

(b) $\Rightarrow$ (a): For estimating derivatives of order at least 1, simply set $V = \Omega$, $\varphi = \text{id}_\Omega$, $L = K$, $W = \Omega'$, $\psi = \text{id}_{\Omega'}$ and $L' = K'$ where K, K' are as in the definition of $\mathcal{E}[\Omega, \Omega']$. $\square$

Based on this local characterization we now proceed to introduce the concept of moderate mappings between manifolds X and Y .

2.2 Definition. *The space $\mathcal{E}_M[X, Y]$ of compactly bounded (c -bounded) moderate maps from X to Y is defined as the set of all $(u_\varepsilon)_\varepsilon \in C^\infty(X, Y)^I$ such that*

$$(i) \quad \forall K \subset\subset \Omega \quad \exists \varepsilon_0 > 0 \quad \exists K' \subset\subset Y \quad \forall \varepsilon < \varepsilon_0 : u_\varepsilon(K) \subseteq K'.$$

$$(ii) \quad \forall k \in \mathbb{N}, \text{ for each chart } (V, \varphi) \text{ in } X, \text{ each chart } (W, \psi) \text{ in } Y, \text{ each } L \subset\subset V \text{ and each } L' \subset\subset W \text{ there exists } N \in \mathbb{N} \text{ with}$$

$$\sup_{x \in L \cap u_\varepsilon^{-1}(L')} \|D^{(k)}(\psi \circ u_\varepsilon \circ \varphi^{-1})(\varphi(p))\| = O(\varepsilon^{-N}).$$

2.3 Remark. (i) Membership of $(u_\varepsilon)_\varepsilon$ in $\mathcal{E}_M[X, Y]$ can equivalently be determined by requiring (ii) merely for charts from given atlases $\mathcal{A}_X = \{(V_\alpha, \varphi_\alpha) \mid \alpha \in A\}$ of X and $\mathcal{A}_Y = \{(W_\beta, \psi_\beta) \mid \beta \in B\}$ of Y . Indeed, suppose that (ii) holds for all charts from $\mathcal{A}_X, \mathcal{A}_Y$ and let $k \in \mathbb{N}$, $L \subset\subset V$, $L' \subset\subset W$, with $(V, \varphi), (W, \psi)$ charts in X and Y , respectively. Choose $V_{\alpha_1}, \dots, V_{\alpha_r}$ covering L and $W_{\beta_1}, \dots, W_{\beta_s}$ covering L' . We may write $L = \bigcup_{i=1}^r L_i$, $L' = \bigcup_{j=1}^s L'_j$ with $L_i \subset\subset V_{\alpha_i}$, $L'_j \subset\subset W_{\beta_j}$. Then for each pair i, j we obtain constants N_{ij} , ε_{ij} and C_{ij} such that the corresponding estimate is satisfied on $L_{\alpha_i} \cap u_\varepsilon^{-1}L'_{\beta_j}$, for all $\varepsilon < \varepsilon_{ij}$. Setting $N := \max N_{ij}$ and $\varepsilon_1 := \min \varepsilon_{ij}$, let $\varepsilon < \varepsilon_1$ and $p \in L$ such that $u_\varepsilon(p) \in L'$. Suppose that $p \in L_i$ and $u_\varepsilon(p) \in L'_j$. Then in some neighborhood of p , $\psi \circ u_\varepsilon \circ \varphi^{-1} = \psi \circ \psi_{\beta_j}^{-1} \circ (\psi_{\beta_j} \circ u_\varepsilon \circ \varphi_{\alpha_i}^{-1}) \circ \varphi_{\alpha_i} \circ \varphi^{-1}$ and the desired estimate follows from the fact that any derivative of $\psi \circ \psi_{\beta_j}^{-1}$ is uniformly bounded on $\psi_{\beta_j}(L'_j)$ and similar for $\varphi_{\alpha_i} \circ \varphi^{-1}$ on $\varphi(L_i)$.

(ii) Expanding once more the Landau symbol in (ii), we obtain (among others) $\exists N \in \mathbb{N} \dots \exists \varepsilon_1 > 0 \quad \exists C > 0 \quad \forall \varepsilon < \varepsilon_1 \dots \leq C\varepsilon^{-N}$. Let us emphasize that, in a certain sense, N and ε_1 have an intrinsic meaning whereas C depends on the charts under consideration. The second part of this assertion is immediate by inspecting the proof of Proposition 2.1; the first part can be made precise as follows: Given $K \subset\subset \Omega$, l and ε_1 can be chosen independently of $V, \varphi, W, \psi, L, L'$ as long as L is allowed to vary only in K . This can be verified by an argument similar to that in the first part of this remark: Choosing K' according to (i) and covering K and K' by the domains V_i resp. W_j of (finitely many) suitable charts, K and K' can be decomposed according to this covering. Each pair i, j giving rise to constants N_{ij} , ε_{ij} , C_{ij} , it is not difficult to show that again $N := \max N_{ij}$ and $\varepsilon_1 := \min \varepsilon_{ij}$ satisfy (the expanded form of) (ii) of Definition 2.2 for all $V, \varphi, W, \psi, L, L'$ as above, provided $L \subseteq K$, in addition.

From Remark 2.3 (i) it follows that $(u_\varepsilon)_\varepsilon \in \mathcal{C}^\infty(X, Y)^I$ is in $\mathcal{E}_M[X, Y]$ if and only if $(u_\varepsilon|_V)_\varepsilon \in \mathcal{E}_M(V, Y)$ for each $V \in \mathcal{V}$ where $\mathcal{V}$ is any open cover of X .

2.4 Definition. Two elements $(u_\varepsilon)_\varepsilon, (v_\varepsilon)_\varepsilon$ of $\mathcal{E}_M[X, Y]$ are called equivalent, $(u_\varepsilon)_\varepsilon \sim (v_\varepsilon)_\varepsilon$, if the following conditions are satisfied:

- (i) For all $K \subset\subset X$, $\sup_{p \in K} d_h(u_\varepsilon(p), v_\varepsilon(p)) \rightarrow 0$ ($\varepsilon \rightarrow 0$) for some (hence every) Riemannian metric h on Y .
- (ii) $\forall k \in \mathbb{N}_0 \forall m \in \mathbb{N}$, for each chart (V, φ) in X , each chart (W, ψ) in Y , each $L \subset\subset V$ and each $L' \subset\subset W$:

$$\sup_{x \in L \cap u_\varepsilon^{-1}(L') \cap v_\varepsilon^{-1}(L')} \|D^{(k)}(\psi \circ u_\varepsilon \circ \varphi^{-1} - \psi \circ v_\varepsilon \circ \varphi^{-1})(\varphi(p))\| = O(\varepsilon^m).$$

That Definition 2.4 (i) is indeed independent of the particular Riemannian metric h follows from Definition 2.2 (i) and Lemma 2 from [12]. To prove that also in Definition 2.4 (ii) it suffices to consider charts from given atlases (and in several instances to follow) we shall make use of the following auxiliary result.

2.5 Lemma. Let $f : \Omega \rightarrow \Omega'$ be continuously differentiable and suppose that $K \subset\subset \Omega$. Then there exists $C > 0$ such that $\|f(x) - f(y)\| \leq C\|x - y\|$ for all $x, y \in K$.

C can be chosen as $C_1 \cdot \sup_{z \in L} (\|f(z)\| + \|Df(z)\|)$ where L is any fixed compact neighborhood of K in Ω and C_1 only depends on L .

Proof. Choose a bump function χ compactly supported in Ω and identical to 1 in a neighborhood of K . Then $\tilde{f} := \chi f$ can be extended to a smooth function on all of $\mathbb{R}^n$ and we may take for C the supremum of $\|D\tilde{f}\|$ on the convex hull of K . The form of C given above follows from the Leibniz rule. $\square$

2.6 Remark. (i) Equivalence of $(u_\varepsilon)_\varepsilon, (v_\varepsilon)_\varepsilon \in \mathcal{E}_M[X, Y]$ can equivalently be determined by requiring (ii) of Definition 2.4 merely for charts from given atlases $\mathcal{A}_X = \{(V_\alpha, \varphi_\alpha) \mid \alpha \in A\}$ of X and $\mathcal{A}_Y = \{(W_\beta, \psi_\beta) \mid \beta \in B\}$ of Y . Indeed, suppose that (ii) holds for all charts from $\mathcal{A}_X, \mathcal{A}_Y$ and let $L \subset\subset V, L' \subset\subset W$, with $(V, \varphi), (W, \psi)$ charts and $k \in \mathbb{N}_0$ and $m \in \mathbb{N}$. Choose $V_{\alpha_1}, \dots, V_{\alpha_r}$ covering K and $W_{\beta_1}, \dots, W_{\beta_t}$ covering L' . We may write $L = \bigcup_{i=1}^r L_i$ and $L' = \bigcup_{j=1}^t L'_j$ with $L_i \subset\subset V_{\alpha_i}, L'_j \subset\subset W_{\beta_j}$. For $1 \leq j \leq t$ let W'_j be a neighborhood of L'_j whose closure is compact and contained in W_{β_j} . By Definition 2.4 (i) there exists ε'_1 such that $\sup_{p \in K} d_h(u_\varepsilon(p), v_\varepsilon(p)) < \min_{1 \leq j \leq t} d_h(L'_j, \partial W'_j)$ for all $\varepsilon < \varepsilon'_1$. Now choose ε''_1 as in (the expanded version of the Landau symbol of) Definition 2.4, (ii) suitable for k, m , and all sets of data $(V_{\alpha_i}, \varphi_{\alpha_i}), (W_{\beta_j}, \psi_{\beta_j}), L_i, \overline{W'_j}$ simultaneously. Set $\varepsilon_1 := \min(\varepsilon'_1, \varepsilon''_1)$. Then if $p \in L$ and $\varepsilon < \varepsilon_1$ are such that $u_\varepsilon(p), v_\varepsilon(p) \in L'$ there exist i such that $p \in L_i$ and j with $u_\varepsilon(p), v_\varepsilon(p) \in W'_j$.

Then in a neighborhood of $x := \varphi(p)$ we have

$$\begin{aligned} & \psi \circ u_\varepsilon \circ \varphi^{-1} - \psi \circ v_\varepsilon \circ \varphi^{-1} \\ &= (\psi \circ \psi_{\beta_j}) \circ (\psi_{\beta_j} \circ u_\varepsilon \circ \varphi_{\alpha_i}) \circ (\varphi_{\alpha_i} \circ \varphi^{-1}) \\ & \quad - (\psi \circ \psi_{\beta_j}) \circ (\psi_{\beta_j} \circ v_\varepsilon \circ \varphi_{\alpha_i}) \circ (\varphi_{\alpha_i} \circ \varphi^{-1}). \end{aligned}$$

Hence the claim follows from the chain rule by taking into account Lemma 2.5.

(ii) Again ε_1 occurring in the expanded version of the Landau symbol in Definition 2.4 (ii) has an intrinsic character whereas the constant C in the respective estimate depends on the charts under consideration. The proof is similar to that of part (i) of this remark, compare (ii) of Remark 2.3.

Similar considerations as in Remark 2.6 show that the relation $\sim$ is transitive. Hence $\sim$ is indeed an equivalence relation. Having introduced the basic building blocks of our construction we may now proceed to defining the space $\mathcal{G}[X, Y]$ itself.

2.7 Definition. *The quotient $\mathcal{G}[X, Y] := \mathcal{E}_M[X, Y] / \sim$ is called the space of compactly bounded (c-bounded) Colombeau generalized functions from X into Y .*

Due to the c-boundedness of representatives, typical singularities that can be modelled by elements of $\mathcal{G}[X, Y]$ are jump discontinuities along hypersurfaces. When differentiating such generalized functions (by means of component-wise tangent maps to be introduced below), however, also delta-type singularities will enter the picture.

2.8 Examples.

- (i) By Proposition 2.1 and a similar result for the respective null ideal of $\mathcal{G}[\Omega, \Omega']$, for $X = \Omega \subseteq \mathbb{R}^n$, $Y = \Omega' \subseteq \mathbb{R}^m$ we obtain precisely $\mathcal{G}[\Omega, \Omega']$ as introduced at the beginning of this section.
- (ii) Let $H = \text{cl}[(H_\varepsilon)_\varepsilon] \in \mathcal{G}(\mathbb{R})$ such that H_ε converges weakly to the Heaviside function. Then $(\exp(i\pi H_\varepsilon))_\varepsilon \in \mathcal{E}_M[\mathbb{R}, S^1]$ may be viewed as a representative of a jump function valued in S^1 (cf. also Example 2.15 (ii) below).
- (iii) Geodesics in generalized spacetimes. Solving geodesic equations in singular spacetimes modelled locally in Colombeau algebras (cf. [11], [19], [20] and the literature cited therein) yields representatives $(u_\varepsilon)_\varepsilon$ where each $u_\varepsilon : J \subseteq \mathbb{R} \rightarrow X$. It can be shown that such “generalized geodesics” are elements of $\mathcal{G}[J, X]$. For details we refer to [13].
- (iv) In [18], R. Penrose introduced a “discontinuous coordinate transformation” T transforming the distributional form

$$ds^2 = f(x)\delta(u)du^2 - du dv + \sum_{i=1}^2 (dx^i)^2$$

of an impulsive gravitational wave into a continuous form. This transformation was analyzed in the context of special Colombeau algebras in [10]. T was recognized as the distributional limit of a family $(T_\varepsilon)_\varepsilon$ of diffeomorphisms. It follows from (30) in [10] that $T = \text{cl}[(T_\varepsilon)_\varepsilon]$ provides an example of an element of $\mathcal{G}[X, X]$.

2.9 Definition. We call two elements $(u_\varepsilon)_\varepsilon, (v_\varepsilon)_\varepsilon$ of $\mathcal{E}_M[X, Y]$ equivalent of order 0, $(u_\varepsilon)_\varepsilon \sim_0 (v_\varepsilon)_\varepsilon$ if they satisfy Definition 2.4 (i) and (ii) for $k = 0$, or, equivalently, (i) and

(ii') $\forall K \subset\subset \Omega \ \forall m \in \mathbb{N} \ \exists \varepsilon_1 > 0 \ \forall (W, \psi) \text{ chart in } Y \ \forall L' \subset\subset W \ \exists C > 0 \text{ such that}$

$$|(\psi \circ u_\varepsilon(p) - \psi \circ v_\varepsilon(p))| \leq C\varepsilon^m$$

for all $\varepsilon < \varepsilon_1$ and all $p \in K$ with $u_\varepsilon(p), v_\varepsilon(p) \in L'$.

The equivalence of (ii') above and (ii) of Definition 2.4 for $k = 0$ (provided $(u_\varepsilon)_\varepsilon, (v_\varepsilon)_\varepsilon \in \mathcal{E}_M[X, Y]$) can be established by the techniques employed in the proofs of Remarks 2.3 and 2.6. As in Remark 2.6 it follows that it suffices to suppose (ii') for charts from any given atlas $\mathcal{A}_Y$. The following result gives a global characterization of $\sim_0$.

2.10 Theorem. Let $(u_\varepsilon), (v_\varepsilon)_\varepsilon \in \mathcal{E}_M[X, Y]$. The following are equivalent.

- (a) $(u_\varepsilon)_\varepsilon \sim_0 (v_\varepsilon)_\varepsilon$.
- (b) For some (hence every) Riemannian metric h on Y , every $m \in \mathbb{N}$ and every $K \subset\subset X$ we have

$$\sup_{p \in K} d_h(u_\varepsilon(p), v_\varepsilon(p)) = O(\varepsilon^m) \quad (\varepsilon \rightarrow 0).$$

Proof. (b) $\Rightarrow$ (a): (i) from Definition 2.9 (adopted from Definition 2.4) is obvious. Concerning (ii'), let us first suppose that L' is contained in a geodesically convex set W' such that $\overline{W'} \subset\subset W$. Given $m \in \mathbb{N}$, choose $\varepsilon' > 0, C' > 0$ such that $\sup_{p \in K} d_h(u_\varepsilon(p), v_\varepsilon(p)) \leq C'\varepsilon^m$ for $\varepsilon < \varepsilon'$. Then for $\varepsilon < \varepsilon', p \in K$ and $u_\varepsilon(p), v_\varepsilon(p) \in L'$ we have

$$d_h(u_\varepsilon(p), v_\varepsilon(p)) = \int_{a_\varepsilon}^{b_\varepsilon} \|\gamma'_\varepsilon(s)\|_h ds \quad (4)$$

where $\gamma_\varepsilon : [a_\varepsilon, b_\varepsilon] \rightarrow W'$ is the unique geodesic in W' joining $u_\varepsilon(p)$ and $v_\varepsilon(p)$. Since W' is relatively compact in W there exists $C'' > 0$ such that $|\xi| \leq C'' \|T_{\psi(p)}\psi^{-1}(\xi)\|_h$ for all $p \in W'$ and all $\xi \in \mathbb{R}^n$ (with $|\cdot|$, say, the Euclidean norm). Thus

$$\begin{aligned} |\psi(u_\varepsilon(p)) - \psi(v_\varepsilon(p))| &\leq \int_{a_\varepsilon}^{b_\varepsilon} |(\psi \circ \gamma_\varepsilon)'(s)| ds \\ &\leq C'' \int_{a_\varepsilon}^{b_\varepsilon} \|\gamma'_\varepsilon(s)\|_h ds. \end{aligned}$$

From this and (4) the claim follows. To treat the general case where L' is any compact subset of W , let W_q be a geodesically convex open neighborhood of q whose closure is compact and contained in W , for each $q \in L'$. Then there

exist $W_{q_1}, \dots, W_{q_k}$ covering L' and we may write $L' = \bigcup_{i=1}^k L'_i$ with $L'_i \subset\subset W_{q_i}$. Choose $\varepsilon'' > 0$ such that $\sup_{p \in K} d_h(u_\varepsilon(p), v_\varepsilon(p)) < \min_{1 \leq i \leq k} d_h(L'_i, \partial W_{q_i})$ for all $\varepsilon < \varepsilon''$. Then if $p \in K$ and $\varepsilon < \varepsilon''$ are such that $u_\varepsilon(p), v_\varepsilon(p) \in L'$ there exists i with $u_\varepsilon(p), v_\varepsilon(p) \in W_{q_i}$. By the first part of the proof there exist $\varepsilon' > 0$ and $C_i > 0$ ($1 \leq i \leq k$) such that $|\psi \circ u_\varepsilon(p) - \psi \circ v_\varepsilon(p)| \leq C_i \varepsilon^m$ whenever $\varepsilon < \varepsilon'$, $p \in K$ and $u_\varepsilon(p), v_\varepsilon(p) \in W_{q_i}$. We may therefore set $\varepsilon_1 = \min(\varepsilon', \varepsilon'')$ and $C = \max_{1 \leq i \leq k} C_i$ to establish the claim.

(a) $\Rightarrow$ (b): Suppose that (b) is violated. Then for some $K \subset\subset X$ we have

$$\exists m_0 \in \mathbb{N} \ \forall k \in \mathbb{N} \ \exists p_k \in K \ \exists \varepsilon_k < \frac{1}{k} : d_h(u_{\varepsilon_k}(p_k), v_{\varepsilon_k}(p_k)) \geq \varepsilon_k^{m_0}. \quad (5)$$

Using Definition 2.2 (i) and choosing suitable subsequences if necessary we may suppose that $p_k \rightarrow p \in K$, $u_{\varepsilon_k}(p_k) \rightarrow q \in Y$ and $v_{\varepsilon_k}(p_k) \rightarrow q$. Choose a chart (W, ψ) such that $q \in W$. For k sufficiently large we have $u_{\varepsilon_k}(p_k), v_{\varepsilon_k}(p_k) \in L' \subset\subset W$ with $L' = \psi^{-1}(\overline{B_r(\psi(q))})$ ($r > 0$ suitable). Hence also the line connecting $\psi(u_{\varepsilon_k}(p_k))$ and $\psi(v_{\varepsilon_k}(p_k))$ is contained in $\psi(L')$. We now multiply the Euclidean metric on $\psi(W)$ by some cut-off function compactly supported in $\psi(W)$ and identical to 1 in a neighborhood of $\overline{B_r(\psi(q))}$ and extend the pullback of the resulting product under ψ to a Riemannian metric g on Y . Then from Definition 2.9 (ii') we conclude

$$d_g(u_{\varepsilon_k}(p_k), v_{\varepsilon_k}(p_k)) \leq |\psi \circ u_{\varepsilon_k}(p_k) - \psi \circ v_{\varepsilon_k}(p_k)| = O(\varepsilon^r)$$

as $\varepsilon \rightarrow 0$ for large enough k and arbitrary $r \in \mathbb{N}$. But by Lemma 2 of [12] we have $d_h(u_{\varepsilon_k}(p_k), v_{\varepsilon_k}(p_k)) \leq C' d_g(u_{\varepsilon_k}(p_k), v_{\varepsilon_k}(p_k))$ for some $C' > 0$ and all k , so we arrive at a contradiction to (5). $\square$

2.11 Remark. It is a rather recent development in the theory of algebras of generalized functions on open subsets Ω of $\mathbb{R}^n$ that a characterization of the Colombeau-ideal $\mathcal{N}(\Omega)$ as a subspace of $\mathcal{E}_M(\Omega)$ can be given where the asymptotic vanishing of representatives has only to be supposed in the 0-th derivative ([6], Th. 13.1 and the remark following it). In our present setting this implies that for elements $(u_\varepsilon)_\varepsilon, (v_\varepsilon)_\varepsilon$ of $\mathcal{E}_M[\Omega, \Omega']$ in fact $(u_\varepsilon)_\varepsilon \sim (v_\varepsilon)_\varepsilon$ is equivalent with $(u_\varepsilon)_\varepsilon \sim_0 (v_\varepsilon)_\varepsilon$. It would certainly be desirable to obtain such an equivalence also in the general case, i.e., for $\mathcal{G}[X, Y]$. However, the Taylor argument used in the local case (based in turn on Landau's paper [14]) is not applicable in this situation. In fact, setting $f_\varepsilon := \psi \circ u_\varepsilon \circ \varphi^{-1} - \psi \circ v_\varepsilon \circ \varphi^{-1}$ the domain D_ε of definition of f_ε may fail to contain $x + \varepsilon^{m+N} e_i$ for any $x \in D_\varepsilon$, so an expression of the form $f_\varepsilon(x + \varepsilon^{m+N} e_i)$ (as required for the proof of [6], Th. 13.1 may be undefined. This “minimal size” of the domain of definition of f_ε , however, is a necessary requirement for estimating $D^{(k)} f_\varepsilon$ by means of $D^{(k-1)} f_\varepsilon$ and $D^{(k+1)} f_\varepsilon$ (cf. [14], Satz 3). Hence there does not seem to be a straightforward adaptation of the $\mathcal{G}[\Omega, \Omega']$ -argument to the manifold setting and we have to leave this as an open problem. See, however, Remark 2.18 below.

Let us now study the action of elements of $\mathcal{G}[X, Y]$ on generalized points. To this end we introduce the following terminology. Let $f : (X, g) \rightarrow (Y, h)$ be a

smooth map between Riemannian manifolds. For $p \in X$ we denote by $\|T_p f\|_{g,h}$ the norm of the linear map $T_p f : (T_p X, \|\cdot\|_g) \rightarrow (T_{f(p)} X, \|\cdot\|_h)$. Then we have

2.12 Lemma. *Let $(u_\varepsilon)_\varepsilon \in \mathcal{E}_M[X, Y]$, g a Riemannian metric on X and h a Riemannian metric on Y . Then for any $K \subset\subset X$ there exist $\varepsilon_0 > 0$, $N \in \mathbb{N}$ and $C > 0$ such that*

$$\sup_{p \in K} \|T_p u_\varepsilon\|_{g,h} \leq C\varepsilon^{-N}$$

for each $\varepsilon < \varepsilon_0$.

Proof. Immediate from the local expression of $\|T_p u_\varepsilon\|_{g,h}$ and the definition of moderateness. $\square$

2.13 Proposition. *Let $u = \text{cl}[(u_\varepsilon)_\varepsilon] \in \mathcal{G}[X, Y]$ and $\tilde{p} \in \tilde{X}_c$. Then $u(\tilde{p}) := \text{cl}[(u_\varepsilon(p_\varepsilon))]$ is a well-defined element of $\tilde{Y}_c$.*

Proof. Let $(u_\varepsilon)_\varepsilon \sim (v_\varepsilon)_\varepsilon$ and suppose that $\tilde{p}$ is supported in $K \subset\subset X$. Then in particular $(u_\varepsilon)_\varepsilon \sim_0 (v_\varepsilon)_\varepsilon$, so by Theorem 2.10 for any Riemannian metric h we have $d_h(u_\varepsilon(p_\varepsilon), v_\varepsilon(p_\varepsilon)) \leq \sup_{p \in K} d_h(u_\varepsilon(p), v_\varepsilon(p)) = O(\varepsilon^m)$ for each $m \in \mathbb{N}$, so $u_\varepsilon(p_\varepsilon) \sim v_\varepsilon(p_\varepsilon)$.

Moreover, we have to show that $(p_\varepsilon)_\varepsilon \sim (p'_\varepsilon)_\varepsilon$ entails $(u_\varepsilon(p_\varepsilon))_\varepsilon \sim (u_\varepsilon(p'_\varepsilon))_\varepsilon$. We first note that for ε sufficiently small, $p_\varepsilon, p'_\varepsilon$ remain in some $K \subset\subset X$. K can be written in the form $\bigcup_{i=1}^s K_i$ with $K_i \subset\subset C_i$, C_i a connected component of X ($1 \leq i \leq s$). By the theorem of Nomizu-Ozeki (see, e.g., [8]) we may equip X with a geodesically complete Riemannian metric g . For the distance d_g induced by g on X we agree upon setting $d_g(p, p') = \infty$ if p and p' do not lie in the same connected component of X . Then for $\varepsilon < \varepsilon_0$ there exists a g -geodesic $\gamma_\varepsilon : [a_\varepsilon, b_\varepsilon] \rightarrow X$ such that

$$d_g(p_\varepsilon, p'_\varepsilon) = \int_{a_\varepsilon}^{b_\varepsilon} \|\gamma'_\varepsilon(s)\|_g ds$$

(where we have used that for ε small, p_ε and p'_ε necessarily lie in the same C_i with $1 \leq i \leq s$ depending on ε). Choosing $r_i \in \mathbb{R}_+$, $p_i \in K_i$ such that $\{p \in X \mid d_g(p, p_i) \leq r_i\} \supseteq K_i$ it follows that each $\gamma_\varepsilon([a_\varepsilon, b_\varepsilon])$ is contained in $K' := \bigcup_{i=1}^s \{p \in X \mid d_g(p, K_i) \leq 2r\}$ with $r = \max_{1 \leq i \leq s} r_i$. K' is compact by the Hopf-Rinow theorem and our above assumption on d_g . Thus since $(u_\varepsilon)_\varepsilon$ is moderate, by Lemma 2.12 there exists $N \in \mathbb{N}$ and for any Riemannian metric h on Y there exists $C > 0$ such that

$$\begin{aligned} d_h(u_\varepsilon(p_\varepsilon), u_\varepsilon(p'_\varepsilon)) &\leq \int_{a_\varepsilon}^{b_\varepsilon} \|(u_\varepsilon \circ \gamma_\varepsilon)'(s)\|_h ds \\ &\leq \int_{a_\varepsilon}^{b_\varepsilon} \|T_{\gamma_\varepsilon(s)} u_\varepsilon\|_{g,h} \|\gamma'_\varepsilon(s)\|_g ds \\ &\leq C\varepsilon^{-N} \int_{a_\varepsilon}^{b_\varepsilon} \|\gamma'_\varepsilon(s)\|_g ds = C\varepsilon^{-N} d_g(p_\varepsilon, p'_\varepsilon). \end{aligned}$$

Since $d_g(p_\varepsilon, p'_\varepsilon) = O(\varepsilon^m)$ for each $m > 0$, it follows that $(u_\varepsilon(p_\varepsilon))_\varepsilon \sim (u_\varepsilon(p'_\varepsilon))_\varepsilon$, as claimed. $\square$

2.14 Proposition. *Let $u = \text{cl}[(u_\varepsilon)_\varepsilon]$, $v = \text{cl}[(v_\varepsilon)_\varepsilon] \in \mathcal{G}[X, Y]$. Then $(u_\varepsilon)_\varepsilon \sim_0 (v_\varepsilon)_\varepsilon$ if and only if $u(\tilde{p}) = v(\tilde{p})$ for all $\tilde{p} \in \tilde{X}_c$.*

Proof. Necessity has already been established in the proof of Proposition 2.13. Conversely, suppose that $(u_\varepsilon)_\varepsilon \not\sim_0 (v_\varepsilon)_\varepsilon$. Then there exist $K \subset\subset X$, $m_0 \in \mathbb{N}$, $\varepsilon_k < \frac{1}{k}$ and $p_k \in K$ such that $d_h(u_{\varepsilon_k}(p_{\varepsilon_k}), v_{\varepsilon_k}(p_{\varepsilon_k})) \geq \varepsilon_k^{m_0}$ for all $k \in \mathbb{N}$. Setting $p_\varepsilon := p_k$ for $\varepsilon_{k+1} < \varepsilon \leq \varepsilon_k$ we obtain an element $\tilde{p} = \text{cl}[(p_\varepsilon)_\varepsilon] \in \tilde{X}_c$ such that $u(\tilde{p}) \neq v(\tilde{p})$. $\square$

The following result demonstrates that composition of generalized functions can be carried out in our present framework. In its formulation (as well as for some results to follow) we shall make use of the following condition. For any $(u_\varepsilon)_\varepsilon \in \mathcal{E}_M[X, Y]$ we consider the property

$$\forall K \subset\subset X \exists \varepsilon_0 > 0 \exists (\psi, W) \text{ chart in } Y \text{ such that } \overline{\bigcup_{\varepsilon < \varepsilon_0} u_\varepsilon(K)} \subset\subset W. \quad (6)$$

It follows from Theorem 2.10 that if any representative $(u_\varepsilon)_\varepsilon$ of some $u \in \mathcal{G}[X, Y]$ satisfies (6) then so does every other. In this case we say that u itself satisfies (6).

2.15 Examples.

(i) (6) is trivially satisfied for each $u \in \mathcal{G}[X, Y]$ in case Y is an open subset of some $\mathbb{R}^m$.

(ii) Let

$$h_\varepsilon(x) := \frac{1+\varepsilon}{2} [\tanh(x) + \tanh(\frac{x}{\varepsilon})]$$

and $u_\varepsilon(x) := e^{i\pi h_\varepsilon(x)}$. Then $u = \text{cl}[(u_\varepsilon)_\varepsilon] \in \mathcal{G}[\mathbb{R}, S^1]$ may be viewed as modelling a function displaying a jump at $x = 0$ from $-i \in S^1$ to $i \in S^1$. Each u_ε is onto S^1 , so no single chart of S^1 can contain $\bigcup_{\varepsilon < \varepsilon_0} u_\varepsilon(\mathbb{R})$ for any $\varepsilon_0 > 0$. Nevertheless u satisfies (6).

2.16 Theorem. *Let $u = \text{cl}[(u_\varepsilon)_\varepsilon] \in \mathcal{G}[X, Y]$, $v = \text{cl}[(v_\varepsilon)_\varepsilon] \in \mathcal{G}[Y, Z]$. Suppose that $(v_\varepsilon)_\varepsilon$ satisfies (6). Then $v \circ u := \text{cl}[(v_\varepsilon \circ u_\varepsilon)_\varepsilon]$ is a well-defined element of $\mathcal{G}[X, Z]$.*

Proof. For each $K \subset\subset X$ there exist $K' \subset\subset Y$ and $\varepsilon'_0 > 0$ such that $u_\varepsilon(K) \subseteq K'$ and there exist $K'' \subset\subset Z$ and $\varepsilon''_0 > 0$ with $v_\varepsilon(K') \subseteq K''$ for $\varepsilon < \varepsilon''_0$. Hence $v_\varepsilon(u_\varepsilon(K)) \subseteq K''$ for $\varepsilon < \min(\varepsilon'_0, \varepsilon''_0)$. To finish the proof that $(v_\varepsilon \circ u_\varepsilon)_\varepsilon$ is moderate let $k \in \mathbb{N}$, (V, φ) a chart in X , $L \subset\subset V$, (U, ζ) a chart in Z and $L'' \subset\subset U$. Choose ε_0 and L' such that $u_\varepsilon(L) \subseteq L'$ for $\varepsilon < \varepsilon_0$. Cover L' by charts (W_i, ψ_i) ($1 \leq i \leq r$) and write $L' = \bigcup_{i=1}^r L'_i$ with $L'_i \subset\subset W_i$. Choose $\varepsilon_1 (\leq \varepsilon_0)$

and N according to Definition 2.2 simultaneously for all $1 \leq k' \leq k$, for $(u_\varepsilon)_\varepsilon$ with respect to all sets of data $(V, \varphi), (W_i, \psi_i), L, L_i$ and for $(v_\varepsilon)_\varepsilon$ with respect to all data $(W_i, \psi_i), (U, \zeta), L', L''$. Then if $\varepsilon < \varepsilon_1$ and $p \in L$ with $v_\varepsilon(u_\varepsilon(p)) \in L''$ there exists i with $u_\varepsilon(p) \in L_i$. Thus we may write $D^{(k)}(\zeta \circ v_\varepsilon \circ u_\varepsilon \circ \varphi^{-1})(\varphi(p))$ as $D^{(k)}((\zeta \circ v_\varepsilon \circ \psi_i^{-1}) \circ (\psi_i \circ u_\varepsilon \circ \varphi^{-1}))(\varphi(p))$ and the claim follows from the chain rule.

To show that $v \circ u$ is well-defined let $(u_\varepsilon)_\varepsilon \sim (u'_\varepsilon)_\varepsilon$. Then by Proposition 2.14 for any $\tilde{p} = \text{cl}[(p_\varepsilon)_\varepsilon] \in \tilde{X}_c$ we have $(u_\varepsilon(p_\varepsilon))_\varepsilon \sim (u'_\varepsilon(p_\varepsilon))_\varepsilon$ and, consequently, $(v_\varepsilon(u_\varepsilon(p_\varepsilon)))_\varepsilon \sim (v_\varepsilon(u'_\varepsilon(p_\varepsilon)))_\varepsilon$. Thus $(v_\varepsilon \circ u_\varepsilon)_\varepsilon \sim_0 (v_\varepsilon \circ u'_\varepsilon)_\varepsilon$, again by Proposition 2.14. In particular, (i) of Definition 2.4 is satisfied.

To establish (ii) let $k \in \mathbb{N}_0$, (V, φ) a chart in X , (U, ζ) a chart in Z , $L \subset\subset V$, $L'' \subset\subset U$. Choose $L' \subset\subset Y$ such that $u_\varepsilon(L) \cup u'_\varepsilon(L) \subseteq L'$ for ε small. We can write $L' = \bigcup_{i=1}^r L'_i$, with $L'_i \subset\subset \overline{W'_i} \subset\subset W_i$, (W_i, ψ_i) charts in Y and W'_i respective open neighborhoods of L'_i . Choose a chart (U_1, ζ_1) as provided by (6) for $(v_\varepsilon)_\varepsilon$ and the compact set $K' := \bigcup_{i=1}^r \overline{W'_i}$, yielding $K'' := \bigcup_{\varepsilon < \varepsilon_2} v_\varepsilon(K') \subset\subset U_1$ for some positive ε_2 . Now suppose $p \in L$ and $v_\varepsilon(u_\varepsilon(p)), v_\varepsilon(u'_\varepsilon(p)) \in L''$. For each fixed ε sufficiently small we have that $u_\varepsilon(p), u'_\varepsilon(p)$ both are contained in the same W'_i for some i . Hence

$$\begin{aligned} & D^{(k)}(\zeta_1 \circ v_\varepsilon \circ u_\varepsilon \circ \varphi^{-1} - \zeta_1 \circ v_\varepsilon \circ u'_\varepsilon \circ \varphi^{-1})(\varphi(p)) \\ &= D^{(k)}((\zeta_1 \circ v_\varepsilon \circ \psi_i^{-1}) \circ (\psi_i \circ u_\varepsilon \circ \varphi^{-1}) \\ &\quad - (\zeta_1 \circ v_\varepsilon \circ \psi_i^{-1}) \circ (\psi_i \circ u'_\varepsilon \circ \varphi^{-1}))(\varphi(p)). \end{aligned}$$

This last expression can be estimated using the chain rule, moderateness of $(v_\varepsilon)_\varepsilon$, equivalence of $(u_\varepsilon)_\varepsilon, (u'_\varepsilon)_\varepsilon$, Lemma 2.5 and moderateness of $(u'_\varepsilon)_\varepsilon$. Note that for each i , Lemma 2.5 is to be applied to the compact subset $\psi_i(\overline{W'_i})$ of the domain of definition $\psi_i(W_i \cap v_\varepsilon^{-1}(U_1))$ of $\zeta_1 \circ v_\varepsilon \circ \psi_i^{-1}$ thus also the constants produced by this part of the argument are in fact independent of ε . Since all derivatives of $\zeta \circ \zeta_1^{-1}$ are bounded on the compact subset $\zeta_1(L'' \cap K'')$ of $\zeta_1(U \cap U_1)$ analogous estimates follow (involving one more application of Lemma 2.5) with ζ replacing ζ_1 in the above. Hence $(v_\varepsilon \circ u_\varepsilon)_\varepsilon \sim (v_\varepsilon \circ u'_\varepsilon)_\varepsilon$.

Finally, if $(v_\varepsilon)_\varepsilon \sim (v'_\varepsilon)_\varepsilon$ then $(v_\varepsilon \circ u_\varepsilon)_\varepsilon \sim_0 (v'_\varepsilon \circ u_\varepsilon)_\varepsilon$ is immediate from property (i) in Definition 2.2 of $(u_\varepsilon)_\varepsilon$. Let $p \in L$ such that $v_\varepsilon(u_\varepsilon(p)), v'_\varepsilon(u_\varepsilon(p)) \in L''$. As above, for small ε we have $u_\varepsilon(p) \in L'_i$, so the estimate for $D^{(k)}((\zeta \circ v_\varepsilon \circ \psi_i^{-1}) \circ (\psi_i \circ u_\varepsilon \circ \varphi^{-1}) - (\zeta \circ v'_\varepsilon \circ \psi_i^{-1}) \circ (\psi_i \circ u_\varepsilon \circ \varphi^{-1}))(\varphi(p))$ follows immediately from the equivalence of $(v_\varepsilon)_\varepsilon$ and $(v'_\varepsilon)_\varepsilon$. $\square$

The map $\sigma : f \mapsto \text{cl}[(f)_\varepsilon]$ provides a canonical embedding of $\mathcal{C}^\infty(Y, Z)$ into $\mathcal{G}[Y, Z]$. Composition with smooth functions can be carried out unrestrictedly:

2.17 Corollary.

- (i) Let $u \in \mathcal{C}^\infty(X, Y)$, $v = \text{cl}[(v_\varepsilon)_\varepsilon] \in \mathcal{G}[Y, Z]$. Then $v \circ u := \text{cl}[(v_\varepsilon \circ u)_\varepsilon]$ is a well-defined element of $\mathcal{G}[X, Z]$.
- (ii) Let $u = \text{cl}[(u_\varepsilon)_\varepsilon] \in \mathcal{G}[X, Y]$, $v \in \mathcal{C}^\infty(Y, Z)$. Then $v \circ u := \text{cl}[(v \circ u_\varepsilon)_\varepsilon]$ is a well-defined element of $\mathcal{G}[X, Z]$.

Proof. The proof of (i) actually is a slimmed-down version of that of Theorem 2.16, with the (hard) part dealing with $(u_\varepsilon)_\varepsilon \sim (u'_\varepsilon)_\varepsilon$ simply omitted. Concerning (ii), we also may proceed along the lines of proof of Theorem 2.16, noting that if $(u_\varepsilon)_\varepsilon \sim (u'_\varepsilon)_\varepsilon$ we may estimate terms of the form

$$\|D^{(k)}((\zeta \circ f \circ \psi^{-1}) \circ (\psi \circ u_\varepsilon \circ \varphi^{-1}) - (\zeta \circ f \circ \psi^{-1}) \circ (\psi \circ u'_\varepsilon \circ \varphi^{-1}))(\varphi(p))\|$$

directly by Lemma 2.5 since both $\psi \circ u_\varepsilon \circ \varphi^{-1}$ and $\psi \circ u'_\varepsilon \circ \varphi^{-1}$ map $L(\subset\subset X)$ into a fixed compact subset of the domain of definition of $\zeta \circ f \circ \psi^{-1}$ for ε small. $\square$

2.18 Remark. Condition (6) can also be employed to remedy the domain problem discussed in Remark 2.11. Thus if $(u_\varepsilon)_\varepsilon, (v_\varepsilon)_\varepsilon \in \mathcal{E}_M[X, Y]$ and $(u_\varepsilon)_\varepsilon$ satisfies (6) then $(u_\varepsilon)_\varepsilon \sim_0 (v_\varepsilon)_\varepsilon$ implies $(u_\varepsilon)_\varepsilon \sim (v_\varepsilon)_\varepsilon$. In fact, by Theorem 2.10, (6) in this case is satisfied simultaneously for $(u_\varepsilon)_\varepsilon$ and $(v_\varepsilon)_\varepsilon$, so the Taylor argument indicated in Remark 2.11 yields the claim.

3 Generalized vector bundle homomorphisms

In the present section we introduce the notion of generalized vector bundle homomorphisms. Among others this notion will allow us to treat tangent mappings of elements of $\mathcal{G}[X, Y]$.

3.1 Definition. Let $\mathcal{E}_M^{\text{VB}}[E, F]$ be the set of all nets $(u_\varepsilon)_\varepsilon \in \text{Hom}(E, F)^I$ satisfying

- (i) $(\underline{u}_\varepsilon)_\varepsilon \in \mathcal{E}_M[X, Y]$.
- (ii) $\forall k \in \mathbb{N}_0 \forall (V, \Phi)$ vector bundle chart in E , $\forall (W, \Psi)$ vector bundle chart in F , $\forall L \subset\subset V \forall L' \subset\subset W \exists N \in \mathbb{N} \exists \varepsilon_1 > 0 \exists C > 0$ with

$$\|D^{(k)}(u_{\varepsilon\Psi\Phi}^{(2)}(\varphi(p)))\| \leq C\varepsilon^{-N}$$

for all $\varepsilon < \varepsilon_1$ and all $p \in L \cap \underline{u}_\varepsilon^{-1}(L')$. Here $\|\cdot\|$ denotes any matrix norm.

Again it suffices to require (ii) merely for charts from arbitrary vector bundle atlases of E resp. F . The proof is entirely analogous to that of Remark 2.3, taking into account the general form (2) of local vector bundle homomorphisms.

3.2 Definition. $(u_\varepsilon)_\varepsilon, (v_\varepsilon)_\varepsilon \in \mathcal{E}_M^{\text{VB}}[E, F]$ are called *vb-equivalent*, denoted by $(u_\varepsilon)_\varepsilon \sim_{vb} (v_\varepsilon)_\varepsilon$, if

- (i) $(\underline{u}_\varepsilon)_\varepsilon \sim (\underline{v}_\varepsilon)_\varepsilon$ in $\mathcal{E}_M[X, Y]$.
- (ii) $\forall k \in \mathbb{N}_0 \forall m \in \mathbb{N} \forall (V, \Phi)$ vector bundle chart in E , $\forall (W, \Psi)$ vector bundle chart in F , $\forall L \subset\subset V \forall L' \subset\subset W \exists \varepsilon_1 > 0 \exists C > 0$ such that:

$$\|D^{(k)}(u_{\varepsilon\Psi\Phi}^{(2)} - v_{\varepsilon\Psi\Phi}^{(2)})(\varphi(p))\| \leq C\varepsilon^m$$

for all $\varepsilon < \varepsilon_1$ and all $p \in L \cap \underline{u}_\varepsilon^{-1}(L') \cap \underline{v}_\varepsilon^{-1}(L')$.

$(u_\varepsilon)_\varepsilon, (v_\varepsilon)_\varepsilon \in \mathcal{E}_M^{\text{VB}}[E, F]$ are called *vb-0-equivalent*, denoted by $(u_\varepsilon)_\varepsilon \sim_{vb} (v_\varepsilon)_\varepsilon$, if $(\underline{u}_\varepsilon)_\varepsilon \sim_0 (\underline{v}_\varepsilon)_\varepsilon$ and (ii) above holds for $k = 0$. We set $\text{Hom}_\mathcal{G}[E, F] := \mathcal{E}_M^{\text{VB}}[E, F] / \sim_{vb}$.

To see that condition (ii) of Definition 3.2 is well behaved under changing vector bundle charts (and thereby that (ii) only has to be required for arbitrary vector bundle atlases) we may again proceed analogously to Remark 2.6. Thus we only have to consider the following local situation: let g, h be local vector bundle homomorphisms, abbreviate $u_{\varepsilon\Psi\Phi}$ by u_ε (and similar for v_ε) and note that $(h \circ u_\varepsilon \circ g)^{(2)} = (x, v) \rightarrow h^{(2)}(u_\varepsilon^{(1)}(g^{(1)}(x)))u_\varepsilon^{(2)}(g^{(1)}(x))g^{(2)}(x)v$. Hence

$$\begin{aligned} & \| (h \circ u_\varepsilon \circ g)^{(2)}(x) - (h \circ v_\varepsilon \circ g)^{(2)}(x) \| \\ & \leq \| h^{(2)}(u_\varepsilon^{(1)}(g^{(1)}(x))) - h^{(2)}(v_\varepsilon^{(1)}(g^{(1)}(x))) \| \| u_\varepsilon^{(2)}(g^{(1)}(x))g^{(2)}(x) \| \\ & \quad + \| h^{(2)}(v_\varepsilon^{(1)}(g^{(1)}(x))) \| \| u_\varepsilon^{(2)}(g^{(1)}(x)) - v_\varepsilon^{(2)}(g^{(1)}(x)) \| \| g^{(2)}(x) \|. \end{aligned}$$

Here the first summand can be estimated using Lemma 2.5 while an estimate of the second one follows immediately from moderateness of $(v_\varepsilon)_\varepsilon$ and Definition 3.2 for $u_\varepsilon^{(2)}, v_\varepsilon^{(2)}$. Derivatives can be treated analogously.

In the case of local vector bundle homomorphisms $u_\varepsilon, v_\varepsilon : \Omega \times \mathbb{R}^{n'} \rightarrow \Omega' \times \mathbb{R}^{m'}$ it follows that $(u_\varepsilon)_\varepsilon \in \mathcal{E}_M^{\text{VB}}(\Omega \times \mathbb{R}^{n'}, \Omega' \times \mathbb{R}^{m'})$ if and only if $(u_\varepsilon^{(1)})_\varepsilon \in \mathcal{E}_M[\Omega, \Omega']$ and $(u_\varepsilon^{(2)})_\varepsilon$ is moderate as a map from Ω into $\mathbb{R}^{m'^2}$ and $(u_\varepsilon)_\varepsilon \sim_{vb} (v_\varepsilon)_\varepsilon$ if and only if $u_\varepsilon^{(i)} - v_\varepsilon^{(i)}$ is negligible in the local sense ($i = 1, 2$).

For $u \in \text{Hom}_\mathcal{G}[E, F]$ the definition $\underline{u} := \text{cl}[(u_\varepsilon)_\varepsilon]$ yields a well-defined element of $\mathcal{G}[X, Y]$ which is uniquely characterized by the property $\underline{u} \circ \pi_X = \pi_Y \circ u$ (with π_X, π_Y the bundle projections).

3.3 Definition. For any $u \in \mathcal{G}[X, Y]$ we define the *tangent map*

$$Tu := \text{cl}[(Tu_\varepsilon)_\varepsilon] \in \text{Hom}_\mathcal{G}(TX, TY).$$

That Tu is well-defined, i.e., that $(u_\varepsilon)_\varepsilon \sim (v_\varepsilon)_\varepsilon$ entails $(Tu_\varepsilon)_\varepsilon \sim_{vb} (Tv_\varepsilon)_\varepsilon$ is immediate from the definitions of $\sim$ and $\sim_{vb}$.

3.4 Definition. On any vector bundle (E, X, π) we consider the set of $(e_\varepsilon)_\varepsilon \in E^I$ such that

- (i) $\exists K \subset\subset X \exists \varepsilon_0 > 0$ such that $(\pi(e_\varepsilon))_\varepsilon \in K$ for all $\varepsilon < \varepsilon_0$.
- (ii) For some Riemannian metric h on E inducing the norm $\| \cdot \|_h$ on the fibers of E we have: $\exists \varepsilon_1 > 0 \exists N \in \mathbb{N} \exists C > 0$ such that

$$\|e_\varepsilon\|_h \leq C\varepsilon^{-N}$$

for all $\varepsilon < \varepsilon_1$. (A different choice of h in this condition only influences C .)

On this set of vb-moderate generalized points we introduce an equivalence relation $\sim_{vb}$ by calling two elements $(e_\varepsilon)_\varepsilon, (e'_\varepsilon)_\varepsilon$ equivalent if

(iii) $(\pi(e_\varepsilon))_\varepsilon \sim (\pi(e'_\varepsilon))_\varepsilon$ in X^I (i.e., $d_g(p_\varepsilon, q_\varepsilon) = O(\varepsilon^s)$ for each s and one (hence every) Riemannian metric g on X).

(iv) $\forall m \in \mathbb{N} \forall (W, \Psi)$ vector bundle chart in $E \forall L' \subset\subset W \exists \varepsilon_1 > 0 \exists C > 0$ such that

$$|\psi e_\varepsilon - \psi e'_\varepsilon| \leq C\varepsilon^m$$

for all $\varepsilon < \varepsilon_1$ whenever both $\pi_X(e_\varepsilon)$ and $\pi_X(e'_\varepsilon)$ lie in L' . (Here $\Psi = (z_p \mapsto (\psi(p), \psi(z)))$, cf. (1).)

The set of equivalence classes is denoted by $E_c^{\sim vb}$.

By Lemma 2.5 it suffices to require (iv) for charts from any given vector bundle atlas. Elements of $E_c^{\sim vb}$ will be written in the form $\tilde{e} = \text{cl}[(e_\varepsilon)_\varepsilon]$. $\pi(\tilde{e}) := \text{cl}[(\pi(e_\varepsilon))_\varepsilon] \in \tilde{X}_c$ is called base point of $\tilde{e}$. For any $\tilde{p} \in \tilde{X}_c$ we set

$$(E_c^{\sim vb})_{\tilde{p}} := \{\tilde{e} \in E_c^{\sim vb} \mid \pi(\tilde{e}) = \tilde{p}\}.$$

Let $\tilde{e} = \text{cl}[(e_\varepsilon)_\varepsilon] \in (E_c^{\sim vb})_{\tilde{p}}$ with $\tilde{p} = \text{cl}[(p_\varepsilon)_\varepsilon]$. Then there exists a representative $(e'_\varepsilon)_\varepsilon$ of $\tilde{e}$ and $\varepsilon_0 > 0$ such that $\pi(e'_\varepsilon) = p_\varepsilon$ for all $\varepsilon < \varepsilon_0$. In fact, there exists $K \subset\subset X$ such that $p_\varepsilon \in K$ for ε small and we may choose vector bundle charts (Ψ_i, V_i) ($1 \leq i \leq k$) such that the V_i cover K . Write $K = \bigcup_{i=1}^k K_i$ with $K_i \subset\subset V_i$ and let V'_i be a neighborhood of K_i whose compact closure is contained in V_i . Then there is $\varepsilon_1 > 0$ such that for every $\varepsilon < \varepsilon_1$ there exists i with $\pi(e_\varepsilon), p_\varepsilon \in \pi(V'_i)$. For $\varepsilon < \varepsilon_1$ we set $e'_\varepsilon = \Psi_i^{-1}(\psi_i(p_\varepsilon), \psi_i(e_\varepsilon))$. Then it follows directly from the definition of $\sim_{vb}$ that $\tilde{e}' = \tilde{e}$, while $\pi(e'_\varepsilon) = p_\varepsilon$ for ε small holds by construction.

This observation paves the way to endowing $(E_c^{\sim vb})_{\tilde{p}}$ with the structure of a $\mathcal{K}$ -module: for $\tilde{e}, \tilde{e}' \in (E_c^{\sim vb})_{\tilde{p}}$ and $\tilde{r} \in \mathcal{K}$ choose representatives $(e_\varepsilon)_\varepsilon, (e'_\varepsilon)_\varepsilon$ as above and define $\tilde{e} + \tilde{r}\tilde{e}' := \text{cl}[(e_\varepsilon + e'_\varepsilon)_\varepsilon]$. Again it follows directly from the definition of $\sim_{vb}$ that $\tilde{e} + \tilde{r}\tilde{e}'$ is well-defined, i.e., it depends exclusively on $\tilde{e}, \tilde{e}'$ and $\tilde{r}$.

Our next aim is to show that $E_c^{\sim vb}$ provides the appropriate concept of point values of generalized sections and generalized vector bundle homomorphisms. To this end we first prove the following strengthening of [12], Lemma 3.

3.5 Lemma. *Let $\tilde{p} = \text{cl}[(p_\varepsilon)_\varepsilon], \tilde{p}' = \text{cl}[(p'_\varepsilon)_\varepsilon] \in \tilde{X}_c$. Then $\tilde{p} = \tilde{p}'$ if and only if*

(i) $d_h(p_\varepsilon, p'_\varepsilon) \rightarrow 0$ for one (each) Riemannian metric h on X .

(ii) For each $m \in \mathbb{N}$, each chart (V, φ) of X and each $L \subset\subset V$ there exist $\varepsilon_1 > 0$ and $C > 0$ with

$$|\varphi(p_\varepsilon) - \varphi(p'_\varepsilon)| \leq C\varepsilon^m$$

whenever $\varepsilon < \varepsilon_1$ and both $p_\varepsilon, p'_\varepsilon$ lie in L .

Proof. This is just an appropriately slimmed-down version of the proof of Theorem 2.10. $\square$

3.6 Proposition.

- (i) Let $u \in \Gamma_{\mathcal{G}}(X, E)$ and $\tilde{p} \in \tilde{X}_c$. Then $u(\tilde{p}) := \text{cl}[(u_{\varepsilon}(p_{\varepsilon}))_{\varepsilon}]$ is a well-defined element of $E_c^{\sim vb}$.
- (ii) Let $v \in \text{Hom}_{\mathcal{G}}[E, F]$ such that $(\underline{v}_{\varepsilon})_{\varepsilon}$ satisfies (6) (for charts on Y induced by vector bundle charts on F) and let $\tilde{e} \in E_c^{\sim vb}$. Then $v(\tilde{e}) := \text{cl}[(v_{\varepsilon}(e_{\varepsilon}))_{\varepsilon}]$ is a well-defined element of $F_c^{\sim vb}$.

Proof. (i) Clearly $(u_{\varepsilon}(p_{\varepsilon}))_{\varepsilon}$ is vb -moderate. To show that $u(\tilde{p})$ is well-defined we first claim that $(u_{\varepsilon})_{\varepsilon} \in \Gamma_{\mathcal{N}}(X, E)$ implies $(u_{\varepsilon}(p_{\varepsilon}))_{\varepsilon} \sim_{vb} (0_{p_{\varepsilon}})_{\varepsilon}$. In fact, (iii) of Definition 3.4 is automatically satisfied and (iv) follows since

$$\|u_{\varepsilon}(p_{\varepsilon})\|_h \leq \sup_{p \in K} \|u_{\varepsilon}(p)\|_h = O(\varepsilon^m)$$

for every $m \in \mathbb{N}$ where $K \subset\subset X$ is such that $p_{\varepsilon} \in K$ for ε small. Supposing now that $(p_{\varepsilon}) \sim (p'_{\varepsilon})$ we have to show that $(u_{\varepsilon}(p_{\varepsilon}))_{\varepsilon} \sim_{vb} (u_{\varepsilon}(p'_{\varepsilon}))_{\varepsilon}$. In this case (iii) of Definition 3.4 holds by assumption. To prove property (iv), cover K by vector bundle charts (V_i, Φ_i) ($1 \leq i \leq k$) such that $K = \bigcup_{i=1}^k K_i$, $K_i \subset\subset V'_i \subset\subset V_i$ and such that in addition $\varphi_i(V'_i)$ is convex for each i . There exists $\varepsilon_1 > 0$ such that for each $\varepsilon < \varepsilon_1$ there is i with $p_{\varepsilon}, p'_{\varepsilon} \in V'_i$. For such ε we have

$$\begin{aligned} & |\varphi_i(u_{\varepsilon}(p_{\varepsilon})) - \varphi_i(u_{\varepsilon}(p'_{\varepsilon}))| \\ &= |\varphi_i \circ u_{\varepsilon} \circ \varphi_i^{-1}(\varphi_i(p_{\varepsilon})) - \varphi_i \circ u_{\varepsilon} \circ \varphi_i^{-1}(\varphi_i(p'_{\varepsilon}))| \\ &\leq \max_{1 \leq i \leq k} \sup_{x \in \varphi_i(V'_i)} \|D(\varphi_i \circ u_{\varepsilon} \circ \varphi_i^{-1})(x)\| |\varphi_i(p_{\varepsilon}) - \varphi_i(p'_{\varepsilon})|. \end{aligned}$$

Here the first factor is bounded by some ε^{-N} since $(u_{\varepsilon})_{\varepsilon}$ is moderate and the second one is $O(\varepsilon^m)$ for each m by Lemma 3.5.

(ii) To show vb -moderateness of $(v_{\varepsilon}(e_{\varepsilon}))_{\varepsilon}$ we first note that $\pi_F(v_{\varepsilon}(e_{\varepsilon}))_{\varepsilon} = (\underline{v}_{\varepsilon}(\pi_X(e_{\varepsilon})))_{\varepsilon}$ is compactly supported, which gives (i) of Definition 3.4. To see Definition 3.4 (ii), choose $K \subset\subset X$ containing $p_{\varepsilon} := \pi_X(e_{\varepsilon})$ for ε small. Then K is covered by finitely many vector bundle charts (V_i, Φ_i) . Writing K as a union of compact sets as above, an estimate of the form

$$|v_{\varepsilon \Psi \Phi_i}^{(2)}(\varphi_i(p_{\varepsilon})) \varphi_i(e_{\varepsilon})| \leq \|v_{\varepsilon \Psi \Phi_i}^{(2)}(\varphi_i(p_{\varepsilon}))\| |\varphi_i(e_{\varepsilon})| \leq \varepsilon^{-N}$$

for some N (using the moderateness assumptions on $(v_{\varepsilon})_{\varepsilon}$ and $(e_{\varepsilon})_{\varepsilon}$) yields the claim.

Suppose that $(e_{\varepsilon})_{\varepsilon} \sim_{vb} (e'_{\varepsilon})_{\varepsilon}$ are supported in K (i.e., $p_{\varepsilon} \in K$ for ε small and analogously for p'_{ε}). We claim that $(v_{\varepsilon}(e_{\varepsilon}))_{\varepsilon} \sim_{vb} (v_{\varepsilon}(e'_{\varepsilon}))_{\varepsilon}$. Property (iii) of Definition 3.4 is immediate from Proposition 2.13, applied to $\underline{v}$. To show (iv) we write $K = \bigcup_{i=1}^r K_i$, $K_i \subset\subset \overline{V'_i} \subset\subset V_i$, (V_i, Φ_i) vector bundle charts in E and V'_i an open neighborhood of K_i . Let $K_1 := \bigcup_{i=1}^r \overline{V'_i} \subset\subset \bigcup_{i=1}^r V_i$ and (using (6)) choose a vector bundle chart (Ψ, W) in Y such that $\overline{\bigcup_{\varepsilon < \varepsilon_1} \underline{v}_{\varepsilon}(K_1)} \subset\subset W$ for some $\varepsilon_1 > 0$. By Definition 3.4 (iii), for each sufficiently small ε there

exists i with $p_\varepsilon, p'_\varepsilon \in V'_i$. Now we note that $(\Psi \circ v_\varepsilon \circ \Phi_i^{-1})(\Phi_i(e_\varepsilon)) = (\psi \circ \underline{v}_\varepsilon \circ \varphi_i^{-1}(\varphi_i(p_\varepsilon)), v_{\varepsilon\Psi\Phi_i}^{(2)}(\varphi_i(p_\varepsilon)) \cdot \varphi_i(e_\varepsilon))$. Hence

$$\begin{aligned} |\psi(v_\varepsilon(e_\varepsilon)) - \psi(v_\varepsilon(e'_\varepsilon))| \\ \leq \|v_{\varepsilon\Psi\Phi_i}^{(2)}(\varphi_i(p_\varepsilon)) - v_{\varepsilon\Psi\Phi_i}^{(2)}(\varphi_i(p'_\varepsilon))\| |\varphi_i(e_\varepsilon)| \\ + \|v_{\varepsilon\Psi\Phi_i}^{(2)}(\varphi_i(p'_\varepsilon))\| |\varphi_i(e_\varepsilon) - \varphi_i(e'_\varepsilon)| \end{aligned}$$

which can be estimated by applying Lemma 2.5. The claim now follows as in the proof of Theorem 2.16 since the constants appearing in this estimate due to the application of Lemma 2.5 are independent of ε : for each i the domain $\varphi_i(V_i \cap \underline{v}_\varepsilon^{-1}(W))$ of $v_{\varepsilon\Psi\Phi_i}^{(2)}$ ($1 \leq i \leq r$) contains the compact set $\varphi_i(\overline{V'_i})$.

Finally, let $(v_\varepsilon)_\varepsilon \sim_{vb} (v'_\varepsilon)_\varepsilon$ (in fact, it suffices to suppose $(v_\varepsilon)_\varepsilon \sim_{vb0} (v'_\varepsilon)_\varepsilon$). Then using the same techniques as above, an estimate of the type

$$|\psi(v_\varepsilon(e_\varepsilon)) - \psi(v'_\varepsilon(e_\varepsilon))| \leq \|v_{\varepsilon\Psi\Phi}^{(2)}(\varphi(p_\varepsilon)) - v'_{\varepsilon\Psi\Phi}^{(2)}(\varphi(p_\varepsilon))\| |\varphi(e_\varepsilon)|$$

yields that $(v_\varepsilon(e_\varepsilon))_\varepsilon \sim_{vb} (v'_\varepsilon(p_\varepsilon))_\varepsilon$. $\square$

Furthermore, a point value characterization based on these concepts is given in the following result.

3.7 Theorem.

- (i) Let $u \in \Gamma_G(X, E)$. Then $u = 0$ if and only if $u(\tilde{p}) = 0_{\tilde{p}}$ in $E_c^{\sim vb}$ for all $\tilde{p} \in \tilde{X}_c$.
- (ii) Let $(v_\varepsilon)_\varepsilon, (v'_\varepsilon)_\varepsilon \in \mathcal{E}_M^{\text{VB}}[E, F]$ as in Proposition 3.6 (ii). Then $(v_\varepsilon)_\varepsilon \sim_{vb0} (v'_\varepsilon)_\varepsilon$ if and only if $v(\tilde{e}) = v'(\tilde{e})$ for all $\tilde{e} \in E_c^{\sim vb}$.

Proof. Necessity is immediate from Proposition 3.6 (i). Conversely, suppose that $u \neq 0$ in $\Gamma_G(X, E)$. Then for some representative $(u_\varepsilon)_\varepsilon$ we have

$$\exists K \subset\subset X \exists m_0 \in \mathbb{N}_0 \forall k \in \mathbb{N} \exists p_k \in K \exists \varepsilon_k < \frac{1}{k} : \|u_{\varepsilon_k}(p_{\varepsilon_k})\|_h > k\varepsilon_k^{m_0}.$$

Setting $p_\varepsilon = p_{\varepsilon_k}$ for $\varepsilon_{k+1} < \varepsilon \leq \varepsilon_k$ we obtain an element $\tilde{p} = \text{cl}[(p_\varepsilon)_\varepsilon]$ of $\tilde{X}_c$ with $u(\tilde{p}) \neq 0_{\tilde{p}}$.

(ii) Necessity follows from the proof of Proposition 3.6 (ii). Conversely, let $(v_\varepsilon)_\varepsilon \not\sim_{vb0} (v'_\varepsilon)_\varepsilon$. Then either $(\underline{u}_\varepsilon)_\varepsilon \not\sim_0 (\underline{v}_\varepsilon)_\varepsilon$ holds or ((ii) for $k = 0$) from Definition 3.2 is violated. In the first case, by Proposition 2.14 there exists $\tilde{p} \in \tilde{X}_c$ with $\underline{v}(\tilde{p}) \neq \underline{v}'(\tilde{p})$. But then $v(0_{\tilde{p}}) \neq v'(0_{\tilde{p}})$. Finally, suppose $\neg((ii) \text{ for } k = 0)$. Then there exist $m \in \mathbb{N}$, vector bundle charts $(V, \Phi), (W, \Psi)$ and $L \subset\subset V, L' \subset\subset W$ allowing the construction of sequences $\varepsilon_j (< 1/j), p_j \in L \cap \underline{v}_{\varepsilon_j}^{-1}(L') \cap \underline{v'}_{\varepsilon_j}^{-1}(L')$ (for $j = 1, 2, \dots$) satisfying

$$\|(v_{\varepsilon_j\Psi\Phi}^{(2)} - v'_{\varepsilon_j\Psi\Phi})(\varphi(p_j))\| > j\varepsilon_j^m.$$

Hence there exists a bounded sequence of vectors $v_j \in \mathbb{R}^{n'}$ such that

$$\|v_{\varepsilon_j}^{(2)}(x_j)v_j - v_{\varepsilon_j}^{\prime(2)}(x_j)v_j\| > j\varepsilon_j^m$$

(with $x_j := \varphi(p_j)$). Now set $e_\varepsilon := \Phi^{-1}(x_j, v_j)$ for $\varepsilon_{j+1} < \varepsilon \leq \varepsilon_j$ to obtain an element $\tilde{e}$ of $E_c^{\sim vb}$ with $v(\tilde{e}) \neq v'(\tilde{e})$. $\square$

Another important transfer process made possible by the above point value concept concerns the alternative description of smooth tensor fields both as sections and multilinear mappings and the resulting character of tensor fields as “pointwise” objects. The following result secures this property also in the context of generalized tensor fields:

3.8 Proposition. *Let $s \in (\mathcal{G})_s^r(X)$ and $\tilde{p} \in \tilde{X}_c$. Let $\omega_1, \dots, \omega_r$ and $\omega'_1, \dots, \omega'_r$ be generalized one-forms with $\omega_i(\tilde{p}) = \omega'_i(\tilde{p})$ ($1 \leq i \leq r$). Let $\xi_1, \dots, \xi_s$ and $\xi'_1, \dots, \xi'_s$ be generalized vector fields with $\xi_i(\tilde{p}) = \xi'_i(\tilde{p})$ ($1 \leq i \leq s$). Then*

$$(s(\omega_1, \dots, \omega_r, \xi_1, \dots, \xi_s))(\tilde{p}) = (s(\omega'_1, \dots, \omega'_r, \xi'_1, \dots, \xi'_s))(\tilde{p}).$$

Proof. As in the proof of the analogous result for smooth tensor fields it suffices to show that for $\omega \in (\mathcal{G})_1^0(X)$ and $\xi \in (\mathcal{G})_0^1(X)$ with $\xi(\tilde{p}) = 0$ we have $(\omega(\xi))(\tilde{p}) = 0$. To begin with, let us suppose that there exists a chart (V, φ) and $K \subset\subset V$ such that $p_\varepsilon \in K$ for ε small.

In the local coordinates (V, φ) we may write $\xi = \sum \xi_j \partial_j$ with $\xi_i \in \mathcal{G}(V)$. Choose a smooth bump function f supported in V such that $f|_K \equiv 1$ in a neighborhood of K . Then $F := \text{cl}[(f)_\varepsilon] \in \mathcal{G}(V)$ and

$$F^2\omega(\xi) = \omega(F^2\xi) = \omega(\sum F\xi_j F\partial_j) = \sum F\xi_j \omega(F\partial_j)$$

from which the result follows upon inserting $\tilde{p}$.

In the general case, let us suppose to the contrary that $\omega(\xi)(\tilde{p}) \neq 0$. Then there exist $m_0 \in \mathbb{N}$, $\varepsilon_k \rightarrow 0$ and representatives $(\omega_\varepsilon)_\varepsilon$, $(\xi_\varepsilon)_\varepsilon$, $(p_\varepsilon)_\varepsilon$ such that

$$|\omega_{\varepsilon_k}(\xi_{\varepsilon_k})(p_{\varepsilon_k})| > \varepsilon_k^{m_0} \quad (7)$$

for all $k \in \mathbb{N}$. Without loss of generality we may suppose that p_{ε_k} converges to some $p \in X$. Choose some chart (V, φ) containing p and some $K \subset\subset V$ such that $p_{\varepsilon_k} \in K$ for large k . Now set $p'_\varepsilon = p_{\varepsilon_k}$ and $\xi'_\varepsilon = \xi_{\varepsilon_k}$ for $\varepsilon_{k+1} \leq \varepsilon \leq \varepsilon_k$. Then we obtain representatives of elements ξ'_ε of $(\mathcal{G})_0^1(X)$ and $\tilde{p}'$ of $\tilde{X}_c$ with $p'_\varepsilon \in K$ for ε small such that $\xi'(\tilde{p}') = 0$ but $\omega(\xi)(\tilde{p}) \neq 0$ by (7), a contradiction to what we have already proved in the special case above. $\square$

Acknowledgments. I am greatly indebted to Michael Grosser for several suggestions that importantly contributed to the final form of the paper. Also, I would like to thank Roland Steinbauer and Andreas Kriegel for helpful discussions.

References

- [1] Aragona, J., Biagioni, H. A. Intrinsic definition of the Colombeau algebra of generalized functions. *Analysis Mathematica*, **17**:75–132, 1991.
- [2] Colombeau, J. F. *New Generalized Functions and Multiplication of Distributions*. North Holland, Amsterdam, 1984.
- [3] Colombeau, J. F. *Elementary Introduction to New Generalized Functions*. North Holland, Amsterdam, 1985.
- [4] Dapić, N., Kunzinger, M., Pilipović, S. Symmetry group analysis of weak solutions. *Proc. London Math. Soc.*, to appear, 2001.
- [5] De Roeper, J. W., Damsma, M. Colombeau algebras on a C^∞ -manifold. *Indag. Mathem., N.S.*, **2**(3), 1991.
- [6] Grosser, M., Farkas, E., Kunzinger, M., Steinbauer, R. On the foundations of nonlinear generalized functions I, II. *Mem. Amer. Math. Soc.* **153**, 2001.
- [7] Grosser, M., Kunzinger, M., Steinbauer, R., Vickers, J. A global theory of algebras of generalized functions. *Adv. Math.*, to appear, 2001.
- [8] Hicks, N. J. *Notes on Differential Geometry*. Number 3 in van Nostrand Mathematical Studies. van Nostrand, New York, 1965.
- [9] Kunzinger, M., Oberguggenberger, M. Group analysis of differential equations and generalized functions. *SIAM J. Math. Anal.*, **31**(6):1192–1213, 2000.
- [10] Kunzinger, M., Steinbauer, R. A note on the Penrose junction conditions. *Class. Quant. Grav.*, **16**:1255–1264, 1999.
- [11] Kunzinger, M., Steinbauer, R. A rigorous solution concept for geodesic and geodesic deviation equations in impulsive gravitational waves. *J. Math. Phys.*, **40**:1479–1489, 1999.
- [12] Kunzinger, M., Steinbauer, R. Nonlinear distributional geometry. *Submitted* (available electronically at <http://arxiv.org/abs/math.FA/0102019>), 2001.
- [13] Kunzinger, M., Steinbauer, R. Generalized pseudo-Riemannian geometry. *Preprint*, 2001.
- [14] Landau, E. Einige Ungleichungen für zweimal differentierbare Funktionen. *Proc. London Math. Soc. Ser. 2*, **13**:43–49, 1913–1914.
- [15] Marsden, J. E. Generalized Hamiltonian mechanics. *Arch. Rat. Mech. Anal.*, **28**(4):323–361, 1968.

- [16] Oberguggenberger, M. *Multiplication of Distributions and Applications to Partial Differential Equations*, volume **259** of *Pitman Research Notes in Mathematics*. Longman, Harlow, U.K., 1992.
- [17] Oberguggenberger, M., Kunzinger, M. Characterization of Colombeau generalized functions by their pointvalues. *Math. Nachr.*, **203**:147–157, 1999.
- [18] Penrose, R. The geometry of impulsive gravitational waves. In L. O’Raifeartaigh, editor, *General Relativity, Papers in Honour of J. L. Synge*, pages 101–115. Clarendon Press, Oxford, 1972.
- [19] Vickers, J. A. Nonlinear generalized functions in general relativity. In Grosser, M., Hörmann, G., Kunzinger, M., Oberguggenberger, M., editor, *Nonlinear Theory of Generalized Functions*, volume **401** of *Chapman & Hall/CRC Research Notes in Mathematics*, pages 275–290, Boca Raton, 1999. CRC Press.
- [20] Vickers, J. A., Wilson, J. P. Generalized hyperbolicity in conical space-times. *Class. Quantum. Grav.*, **17**:1333–1360, 2000.